

Vegetation plays an important role in absorbing carbon dioxide and accelerating the achievement of carbon neutrality. As the ecological barrier of North China, the Taihang Mountains are pivotal to the ecological construction project of China. Nevertheless, the dynamic development of the vegetation carbon sink in the region and the impact factors on the sink ...

Tangyin fault depression, southeast of Taihang mountain which runs through Anyang, Hebi, and Xinxiang, Henan province (). It runs north-Fig. 1 northeast along a strip about 100 kilometers long and 10 to 15 kilometers wide, east-west. The field is situated on the Eastern North China Craton (NCC) between Taihang mountain and the Bohai basin. It

The EE intensity had the strongest explanatory power for the SCSs in the hilly zone, explaining 34.63%. ... In this study, the soil conservation services (SCSs) from 1980 to 2020 in the Taihang Mountain area was assessed using the integrated valuation of ecosystem services and trade-offs (InVEST) model, and the spatial and temporal ...

Aerial photo taken on November 24, 2018 shows photovoltaic power stations on Taihang Mountain in Handan, Hebei Province. The photovoltaic power sector is being pointed to in China as another way in which less-developed, mountainous regions can both save and generate money through solar power.

The central Taihang Mountain region is divided into three ecological zones: a hilly zone (<500 m), mid-mountain zone (500-1500 m), and sub-alpine zone (>1500 m) [38]. Each zone is characterized by ...

The present study verifies that the early Neoproterozoic TTG gneisses and potassic granites in the southeast base of Taihang Mountain emplaced at 2712 ± 65 to 2644 ± 25 Ma, followed by ~2.57 to 2.50 Ga metamorphism and partial anatexis.

An carbon neutrality industrial chain of "desert-photovoltaic power generation-ecological agriculture": Practice from the Ulan Buh Desert, Dengkou, Inner Mongolia. China Geology, 5(3), 549-552. doi: 10.31035/cg2022053. Citation: Chen Xi-jie, Jia Li-qiong, Jia Ting, Hao Zi-guo. 2022. An carbon neutrality industrial chain of "desert ...

Large-scale photovoltaic solar panels have been installed on the Taihang Mountains in Shexian county, North China's Hebei province, to make use of large mountainous areas and to promote clean energy. The installed ...

The highway plant slope of the Taihang Mountain highway in the Hebei province was evaluated using the

assessment model after dividing the highway plant slope stability into four grades.

The Taihang Mountain Region (TMR) of China is an important headwater area that provides substantial water resources supporting the social and ecological stability of the North China Plain.

In the east is the Zhongshan Plateau with Taihang Mountain as the main vein and Qinlu Plateau as the main body. ...  $C_0 + C$  represents the base value that represents the maximum variation of ... the total annual potential of solar photovoltaic power generation in the most suitable area, more suitable area and general area is 11,218.26 GWh ...

During this period, the Taihang Mountains (THM) became a prominent geomorphic and ecological separation zone, which separates NC into regions with different landforms, tectonic settings, and climatic regimes. The uplift history of the THM during the late Mesozoic is of great interest, since it is critical to constrain the geodynamic and ...

Furthermore, there are uncertain factors such as solar radiation intensity, wind speed, runoff and power load in the hybrid system [6], which has caused great difficulties for dispatchers to develop short-term power generation plans. Therefore, the focus of this study is to propose a solution framework for the probabilistic optimal operation model of a WSH hybrid ...

QUYANG, Nov. 29 (Xinhua) -- A solar farm in the Taihang Mountain is generating clean power for Quyang County, north China's Hebei Province. The photovoltaic (PV) power station covers an ...

Numerous studies suggest that various secular geologic and geochemical transitions occurred between ~3.2 Ga and 2.5 Ga. During this age-window, the ~2.70 Ga tectono-thermal event of the Neoproterozoic is by far the most influential, and is unusual in terms of coeval mafic-felsic magmatic rocks which were interpreted to reflect widespread crustal accretion.

The spatial prediction of geothermal sites along the southeast side of Taihang Mountain in the Tangyin rift is a critical goal in the development of renewable energy resources in cities such as ...

Web: <https://arcingenieroslaspalmas.es>